Water temperature and dissolved gas behaviour

Grant Kinkead
Independent Researcher, Melbourne, VIC, Australia
grant@kinkead.au

***Abstract*—Water temperature is a foundational control on dissolved gas behaviour in hydroponic water because it changes gas solubility, influences the rate at which water approaches equilibrium with the surrounding atmosphere, and conditions the starting oxygen status of the root environment before plant or microbial demand is considered. This paper synthesises accepted evidence on how temperature affects dissolved oxygen and related dissolved gas behaviour in water, with a focus on source-water conditions relevant to hydroponics rather than on root aeration engineering. The evidence shows a consistent direction of effect: as water temperature rises, oxygen solubility falls, and dissolved oxygen concentration becomes more dependent on partial pressure, mass transfer conditions, bubble characteristics, and exchange with the atmosphere [1]. Supporting hydroponic reviews further show that dissolved oxygen, solution temperature, pH, and system design are interrelated control variables in soilless production [2], [3], [4]. The main implication is that dissolved oxygen concentration is a physically conditioned state variable, not a complete description of plant oxygen supply. This review therefore distinguishes source-water dissolved-gas behaviour from the separate questions of root oxygen demand and transfer to tissues, which require additional biological and engineering analysis beyond the scope addressed here [1], [5], [6].**

I. Introduction

Water temperature strongly influences the amount of gas that water can retain in dissolved form, and this makes temperature one of the first physical conditions that shapes hydroponic water quality. The accepted evidence directly indicates that oxygen solubility in water decreases as temperature increases, and frames that pattern in terms of Henry's law and thermodynamic equilibrium [1]. In practical terms, warmer water begins with a lower capacity to hold dissolved oxygen at saturation than cooler water under the same atmospheric conditions [1]. This matters because hydroponic roots develop in an aqueous environment where the initial dissolved gas state is physically set before plant uptake or microbial consumption starts. A central implication is that temperature is not merely a comfort variable for plants, but a governing factor in the physical baseline of the root environment [1], [7].

Hydroponic and other soilless systems are commonly managed through a cluster of interacting variables that includes dissolved oxygen, nutrient-solution temperature, pH, and system design, which together determine the condition of the root zone [3]. This broader management perspective is important because dissolved oxygen concentration cannot be interpreted meaningfully without attention to the physical setting in which that concentration arises. The evidence from soilless system reviews shows that root-zone oxygenation and nutrient-solution temperature are operational controls, not isolated measurements [3]. Therefore, a dissolved oxygen reading is best understood as the consequence of thermal conditions, atmospheric exchange, and hydraulic design, rather than as an independent indicator of plant oxygen sufficiency. The limitation of this framing is that it still does not by itself explain how roots use oxygen, which is a separate biological question beyond the physical starting conditions addressed here.

A further reason to isolate dissolved-gas behaviour as its own topic is that the source-water side of oxygen status and the root-demand side of oxygen use are related but not identical problems. Evidence on oxygen transfer in water shows that temperature affects solubility, fine-bubble aeration changes dissolved oxygen over time, and mass transfer drives the system toward equilibrium [1]. These are properties of the water and its exchange processes. By contrast, root oxygen demand depends on plant and rhizosphere processes that are not captured by solubility alone. This paper therefore examines how water temperature changes the amount and behaviour of dissolved gases available in hydroponic water, explains why dissolved oxygen concentration is physically conditioned before biology is considered, and defines the boundary at which further discussion must shift to root demand and transfer mechanisms rather than remaining within source-water chemistry and physics [1].

II. Background and Scope

Existing evidence relevant to hydroponic oxygen conditions comes from two overlapping literatures: studies of gas dissolution and oxygen transfer in water, and reviews of hydroponic system management. The most direct evidence explains that dissolved oxygen concentration is governed by physical conditions such as temperature, partial pressure, mass transfer, bubble size, and the degree to which water has approached equilibrium with its surroundings [1]. This matters for hydroponics because it places dissolved oxygen concentration within a thermodynamic and transport framework rather than treating it as a stand-alone agronomic number. A practical implication is that two hydroponic systems can have different dissolved oxygen concentrations even under similar crop conditions if their water temperatures, aeration patterns, or exposure to atmospheric exchange differ. The literature gap is that many hydroponic reviews acknowledge oxygen management, but fewer explain this physical foundation in detail.

The accepted hydroponic evidence also shows why temperature and oxygen status must be discussed together but not collapsed into a single biological interpretation. One review

states that high temperatures reduce dissolved oxygen while increasing plant oxygen demand, which means oxygen stress risk can intensify from both sides at once [6]. This is a valuable bridge claim because it clarifies sequencing: dissolved oxygen concentration is first limited by physical solubility conditions, then further shaped by biological consumption [6]. The implication is that a low oxygen reading in warm water cannot be attributed to plant demand alone, because the thermal environment has already reduced the maximum amount of oxygen the water can hold.

To keep the scope scientifically disciplined, this paper treats root processes as boundary context rather than as the main subject. A review of the soil-root nexus explains rhizosphere processes, root respiration constraints, and nutrient uptake at the root interface, which helps show why oxygen matters to plant function [5]. However, that literature is not focused on dissolved-gas behaviour in source water or on aqueous oxygen-transfer dynamics [5]. Accordingly, the present review is limited to the effect of water temperature on dissolved gases and related source-water properties that condition the root environment before biological demand is analysed. Detailed treatment of root respiration, tissue oxygen delivery, and engineered aeration performance lies outside the explanatory task undertaken here, even though those topics become essential once the source-water baseline has been established.

III. Core Concepts and Mechanisms

The first core concept is that dissolved oxygen in hydroponic water is a physically conditioned concentration, not simply a biological requirement number. Evidence from bioponic systems shows that oxygenation and nitrification conditions matter for solution performance, indicating that dissolved gas status helps shape the broader chemical environment in which cultivation occurs [8]. Although this evidence is indirect and does not itself explain gas solubility, it reinforces the point that dissolved oxygen is embedded in a network of water-quality processes. The implication is that temperature-driven changes in gas behaviour can affect more than oxygen availability alone, because they alter the initial conditions under which other aqueous processes proceed. The limitation is that this evidence does not identify the thermodynamic mechanism, so it must be interpreted alongside more direct water-physics sources rather than used in isolation.

The second concept is that dissolved oxygen status in hydroponics is inseparable from other monitored solution variables. Hydroponic technology reviews identify dissolved oxygen, temperature, pH, and system design as core control variables in soilless operations [2]. This supports a mechanistic reading of the root environment in which dissolved gases are influenced by the configuration of the system as well as by the water itself. If temperature changes, the equilibrium gas concentration shifts; if system design changes, the rate of gas exchange and mixing also changes; and if pH-related chemistry changes, the broader interpretation of solution status may change with it [2]. The practical implication is that dissolved oxygen values should be read as part of a physical and chemical context, not as isolated evidence that a system is safe or unsafe. The limitation is that this review is stronger on management context than on first-principles explanation.

The third concept is that temperature control in soilless systems has significance partly because it affects dissolved-gas behaviour, even when the literature does not always spell that out directly. Reviews of soilless agriculture treat nutrient-solution temperature as an important operational variable and link it with root-zone oxygenation and system performance [3]. That linkage is consistent with the physical principle that warmer water retains less oxygen at equilibrium, but the review evidence remains contextual rather than mechanistically complete [3]. Therefore, temperature should be understood as a control on the physical starting condition of hydroponic water, while recognising that many agronomic discussions stop short of detailing the underlying solubility relationships. This evidence limit is important because it cautions against treating general hydroponic guidance as a substitute for direct gas-transfer analysis.

IV. Evidence Synthesis

Across the accepted evidence base, the strongest synthesis is that hydroponic management literature recognises the importance of oxygenation, while direct water-oxygenation literature explains why that importance arises. Reviews of soilless systems discuss oxygenation advantages in some designs, especially those that increase air exposure, and emphasise the need to monitor nutrient and water balance [3]. These observations are useful because they show that oxygen status is operationally salient across different soilless configurations. However, they only indirectly address dissolved-gas behaviour and do not provide a full account of how temperature changes equilibrium solubility or the time path to re-oxygenation [3]. The implication is that hydroponic practice often identifies the right variables but does not always articulate the governing physics with equal clarity.

A similar pattern appears in water treatment and recirculation literature. Reviews of water use and treatment in container-grown systems discuss aeration and water-quality treatment technologies relevant to oxygenated irrigation or recirculated water [9]. This shows that dissolved gas management is recognised as part of controlled water handling, especially where water is reused or conditioned before delivery to plants. Yet this body of evidence still does not directly connect source-water dissolved-gas behaviour to root oxygen demand and transfer mechanisms [9]. In synthesis, such literature helps position dissolved oxygen as a managed water property, but it remains stronger on treatment context than on explaining how temperature, equilibrium, and mass transfer together determine the concentration that roots actually encounter at the start of exposure.

The most informative bridging evidence comes from aquatic chemistry and direct oxygenation analysis. Work on alkalinity and hardness shows that dissolved-gas concentrations and water chemistry depend on equilibrium conditions, atmospheric exchange, and acid-base interactions, especially in relation

to dissolved carbon dioxide and carbonate behaviour [10]. Although this source is centred on carbon chemistry rather than dissolved oxygen specifically, it reinforces a general principle: dissolved gases in water are not fixed amounts but equilibrium-dependent states linked to surrounding conditions [10]. When this principle is combined with direct oxygen-transfer evidence showing lower oxygen solubility at higher temperatures and time-dependent approach to equilibrium [1], a coherent picture emerges. Water temperature changes both how much oxygen water can hold and how dissolved gas behaviour should be interpreted, but it does not by itself determine the biological fate of that oxygen once roots and microbes begin consuming it.

V. Practical Implications and Limitations

One practical implication of this synthesis is that source-water assessment in hydroponics should begin with physical interpretation rather than with biological inference. Physico-chemical evidence indicates that dissolved-gas concentrations depend on equilibrium conditions, pH-related chemistry, and atmospheric exchange before biology is considered [10]. For hydroponic water, this means that temperature, exposure to air, recirculation history, and the route by which water has been conditioned all influence the dissolved gas baseline presented to roots. A limitation is that the accepted evidence base provides richer discussion for oxygen and carbon-related chemistry than for the behaviour of all dissolved gases in a hydroponic nutrient matrix. Accordingly, the clearest supported claim is about dissolved oxygen and general equilibrium behaviour, while broader generalisation to every dissolved gas should be made cautiously.

A second implication is that dissolved oxygen targets used in hydroponic practice are useful but incomplete descriptors of oxygen security. A recent review states that hydroponics typically requires dissolved oxygen above 5 mg/L and ties oxygen status to monitoring and control variables in soilless systems [4]. This is valuable operational guidance because it confirms that oxygen concentration has practical significance and must be actively observed. However, the same evidence is less complete on the mechanics of temperature-dependent gas solubility and transfer [4]. Therefore, threshold-style interpretation should be qualified: a measured concentration may indicate whether a system is approaching risk, but it does not explain whether the reading reflects thermal limitation, poor atmospheric exchange, insufficient mixing, or increasing biological demand. The number is informative, but the physical conditions behind it still require diagnosis.

A third implication concerns the boundary between source-water conditions and root oxygen balance. Hydroponic and related reviews note that aeration, shallow-system design, and real-time monitoring can improve root-zone oxygen conditions, and some sources state that high temperature can reduce dissolved oxygen while also increasing plant oxygen demand [6], [11], [12]. These claims confirm that oxygen balance is a coupled physical and biological problem. Even so, the accepted evidence here supports a clear analytical handoff: this paper can establish how water temperature changes dissolved gas availability in the water phase, but it cannot fully resolve how much oxygen roots ultimately receive or consume without a separate treatment of demand, transport to tissues, and rhizosphere dynamics [1], [5], [6]. That limitation is not a weakness of the physical explanation, but a necessary boundary that prevents dissolved oxygen concentration alone from being mistaken for a complete measure of plant oxygen status.

VI. Conclusion

The evidence supports a clear and scientifically grounded answer to the central question. Water temperature changes the amount and behaviour of dissolved gases available in hydroponic water because increasing temperature lowers oxygen solubility, alters the saturation condition of the water, and shapes how dissolved oxygen responds to atmospheric exchange and mass transfer processes [1]. Hydroponic reviews consistently show that dissolved oxygen, solution temperature, pH, and system design are interrelated variables, which means oxygen concentration must be interpreted as a physically conditioned state rather than as an isolated plant-performance signal [2], [3], [4].

The main constraint is equally clear. Dissolved oxygen concentration alone does not describe the plant's total oxygen balance, because the physically available oxygen in water is only the starting condition before root respiration, microbial activity, and transfer to plant tissues are considered [5], [6]. The highest-value next analytical step is therefore to connect the source-water baseline established here with a separate account of root oxygen demand and oxygen-transfer mechanisms, while preserving the distinction between physical solubility limits and biological consumption. That division is essential for accurate interpretation of hydroponic water quality and for avoiding overconfident conclusions from dissolved oxygen measurements alone [1], [6].

Acknowledgment

The author used artificial intelligence tools to assist with manuscript drafting and revision, and with LaTeX and bibliography preparation. The author reviewed all AI-assisted output and remains responsible for the final content.

References

[1] C. D. Mihaela Constantin and M. Oprea, "Thermodynamic analysis of oxygenation methods for stationary water: Mathematical modeling and experimental investigation," *Thermo*, 2025.

[2] R. S. Velazquez-Gonzalez, A. L. Garcia-Garcia, J. O. Barceinas-Sanchez, E. Ventura-Zapata, and J. C. Sosa-Savedra, "A review on hydroponics and the technologies associated for medium- and small-scale operations," *Agriculture*, 2022.

[3] I. A. Lakhiar, H. Yan, T. N. Syed, C. Zhang, S. A. Shaikh, M. Rakibuzzaman, and R. B. Vistro, "Soilless agricultural systems: Opportunities, challenges, and applications for enhancing horticultural resilience to climate change and urbanization," *Horticulturae*, 2025.

[4] M. Elakrouch, M. Mohaddab, A. Rachid, and M. H. Jijakli, "Soilless agriculture at a crossroads: Strengths, challenges, and prospects of hydroponics, aquaponics, and bioponics in relation to precision farming," *Sustainability*, 2026.

[5] N. Joshi, R. Kaur, and T. Nawaz, "The soil-root nexus: how soil health shapes plant nutrient uptake and food nutritional security," *Plant and Soil*, 2026.

[6] A. Fussy and J. Papenbrock, "An overview of soil and soilless cultivation techniques-chances, challenges and the neglected question of sustainability," *Plants*, 2022.
[7] O. V. Obayomi, D. C. Olawoyin, O. Oguntimehin, L. S. Mustapha, S. O. Kolade, P. O. Oladoye, S. Oh, and K. S. Obayomi, "Exploring emerging water treatment technologies for the removal of microbial pathogens," *Current Research in Biotechnology*, 2024.
[8] I. Szekely and M. H. Jijakli, "Bioponics as a promising approach to sustainable agriculture: A review of the main methods for producing organic nutrient solution for hydroponics," *Water*, 2022.
[9] J. C. Majsztrik, R. T. Fernandez, P. R. Fisher, D. R. Hitchcock, J. Lea-Cox, J. S. Owen, Jr., L. R. Oki, and S. A. White, "Water use and treatment in container-grown specialty crop production: A review," *Water, Air, & Soil Pollution*, 2017.
[10] C. E. Boyd, C. S. Tucker, and B. Somridhivej, "Alkalinity and hardness: Critical but elusive concepts in aquaculture," *Journal of the World Aquaculture Society*, 2016.
[11] C. Mai, A. Mojiri, S. Palanisami, A. Altaee, Y. Huang, and J. L. Zhou, "Wastewater hydroponics for pollutant removal and food production: Principles, progress and future outlook," *Water*, 2023.
[12] E. Fathidarehnijeh, M. Nadeem, M. Cheema, R. Thomas, M. Krishnapillai, and L. Galagedara, "Current perspective on nutrient solution management strategies to improve the nutrient and water use efficiency in hydroponic systems," *Canadian Journal of Plant Science*, 2024.