Root respiration in hydroponic systems

Grant Kinkead
Independent Researcher, Melbourne, VIC, Australia
grant@kinkead.au

***Abstract*—Hydroponic cultivation depends on direct management of the root environment, yet oxygen is sometimes treated as a secondary engineering variable rather than a primary biological requirement. The central problem is that roots do not photosynthesise and therefore cannot rely on light exposure to meet their own immediate energy needs. This paper explains why living roots require oxygen for aerobic respiration and how root metabolic demand creates a continuous oxygen requirement in hydroponic root zones. Drawing on plant respiration literature and hydroponic reviews, the paper shows that oxygen supports mitochondrial oxidative phosphorylation, which supplies the ATP needed for maintenance, membrane transport, nutrient uptake, hydraulic function, growth, and defence responses [1], [2]. When oxygen becomes limiting, roots are forced towards fermentative metabolism with much lower ATP yield, creating functional constraints even before visible failure appears [1]. The evidence further indicates that oxygen demand is not static: it rises with metabolically active stages such as establishment and intensive root development, while hydroponic system conditions can either support or restrict oxygen availability [2], [3]. The practical implication is that oxygen provision in hydroponics is not optional optimisation. It is a continuous physiological requirement that must be managed if root function, nutrient acquisition, and crop performance are to be sustained [4], [5].**

I. Introduction

In hydroponic systems, the root environment is engineered directly, so any misunderstanding of root oxygen needs can become a design error rather than a minor cultural issue. Living root tissues require oxygen because aerobic respiration in mitochondria drives oxidative phosphorylation, the process that generates most of the ATP used for cellular work. When oxygen supply falls, this pathway is suppressed and plant tissues are pushed towards fermentative metabolism, which yields far less usable energy and is associated with metabolic stress and adaptive survival responses rather than efficient growth [1]. This makes oxygen a biological requirement of the root zone, not merely a convenience for improving vigour or preventing obvious disorder [6].

The reader question addressed here is why hydroponic roots need oxygen even though the shoot is exposed to light. The key point is that light captured by leaves does not remove the root's immediate dependence on oxygen for its own mitochondrial ATP generation. The evidence shows that oxygen availability is tightly linked to ATP production, energy limitation, and metabolic switching in plant tissues, including roots, so metabolically active root cells still need oxygen to sustain aerobic energy supply [1]. In practical terms, nutrient solution delivery and illumination above the root zone cannot substitute for the oxygen required within root tissues themselves.

This paper develops an evidence based explanation of continuous root oxygen demand in hydroponics. It first defines the physiological scope of root respiration, then examines how metabolic activity influences oxygen demand, and finally considers what current hydroponic literature does and does not establish about system level oxygen management [1]. The central argument is that roots require oxygen continuously because their maintenance, uptake, growth, and protective functions depend on ongoing aerobic respiration, including periods when shoots are not actively photosynthesising or when system conditions restrict oxygen transfer to the root zone [1].

II. Background and Scope

Existing hydroponic literature often treats oxygen as part of general root zone management, but the strongest direct explanation of root respiration in the supplied evidence comes from work linking hypoxia to functional failure in roots. In strawberry nutrient film systems, dissolved oxygen limitation is associated with reduced oxidative phosphorylation, ATP depletion, impaired active nutrient uptake, lower hydraulic conductance, and poorer crop performance [2]. This evidence is important because it connects respiration to concrete root functions rather than to an abstract preference for aeration. Aerobic respiration matters because roots must continuously power transport and maintenance processes that cannot proceed normally when ATP supply collapses.

The same evidence base also clarifies that root oxygen demand is dynamic rather than fixed. Root demand becomes especially critical during establishment and fruit development, and expanding root mats can intensify local oxygen depletion by increasing biological demand within the root zone [2]. This directly supports the relationship between root metabolic activity and oxygen requirement. A hydroponic system that appears adequate under low demand conditions may therefore become limiting as roots enlarge or developmental priorities shift. The practical implication is that oxygen management should be understood in relation to living metabolic load, not only in relation to water movement or nominal system type.

The scope of this paper is therefore explanatory rather than experimental. It does not attempt to derive a universal dissolved oxygen target for all hydroponic crops, because the supplied evidence does not support that level of generalisation. Instead, it explains why roots need oxygen continuously: root ATP production, membrane transport, nutrient uptake, hydraulic conductance, and defence capacity all depend on

oxidative phosphorylation, and these requirements persist regardless of whether shoots are currently supplying photosynthate through active illumination [2]. The gap in much broader hydroponic commentary is that oxygenation is frequently recommended, yet the physiological reason for continuous root demand is often stated only briefly or indirectly.

III. Core Concepts and Mechanisms

A useful starting mechanism is that hydroponic root zones are chemically and biologically dynamic environments. In a nutrient film technique lettuce study, recirculating solution composition changed over time, likely reflecting ongoing biological and chemical transformations, and the authors showed that electrical conductivity alone could miss important root zone imbalances [7]. Although this study does not directly explain continuous oxygen requirement, it reinforces a key conceptual point: root systems are active metabolic interfaces, not passive structures suspended in solution. Because root respiration occurs within this changing environment, oxygen provision cannot be inferred reliably from simplified bulk indicators alone.

Comparative soilless literature helps explain why certain system designs often perform better for oxygen supply without, by itself, resolving the underlying physiology. Reviews of soilless cultivation note that aeroponics can improve root oxygenation because roots are exposed to air while receiving intermittent nutrient delivery, reducing the diffusion constraints typical of continuously water based systems [8]. This supports the idea that oxygen availability influences root performance, yet it remains only a partial explanation of root respiration. Better oxygenation is advantageous not because air exposure is inherently superior, but because root cells still rely on aerobic metabolism to meet their ongoing energy demand.

The mechanism linking design and physiology is therefore conditional rather than purely structural. Aeroponic and other carefully controlled systems may facilitate oxygen transfer, but the relevant biological question is whether oxygen supply keeps pace with root metabolic activity across time and crop stage [8]. Broader environmental control in soilless systems can improve the root environment, yet the supplied review does not directly quantify how changing metabolism sets oxygen demand [8]. That limitation matters because system categories alone do not answer the reader question. Roots need oxygen continuously because respiration is continuous, while system design only changes how easily that need can be met.

IV. Evidence Synthesis

Across the hydroponic reviews in the evidence base, there is broad agreement that oxygenation is important, but the reasons are unevenly developed. One review on medium and small scale hydroponics highlights the importance of oxygenation and maintaining suitable root environments, which is consistent with practical grower guidance, yet it does not directly analyse how metabolic activity within root tissues determines oxygen demand [9]. When this is read alongside the plant respiration literature, the interpretation becomes clearer: oxygen is necessary because roots must sustain mitochondrial ATP production, not simply because oxygenated water is a generic hallmark of good practice [1]. The synthesis value lies in joining hydroponic management statements to the underlying respiratory mechanism.

A similar pattern appears in broader reviews of soilless agriculture. Aeroponics is often described as advantageous because it provides strong root oxygenation, while enclosed soilless systems more generally require careful environmental control to maintain suitable conditions for growth [3]. These observations are compatible with the physiological evidence, but they are still mostly system centred descriptions. Their explanatory power improves when paired with direct evidence that oxygen limitation disrupts oxidative phosphorylation and ATP availability in roots [1], [2]. In other words, the engineering preference for better oxygenation reflects a deeper biological necessity: oxygen supply must support continuous respiratory metabolism in living root tissues.

This synthesis also answers the question of why root oxygen demand persists irrespective of current shoot photosynthesis. Water based and air exposed systems differ in their capacity to maintain root oxygenation, yet those design differences do not alter the root's dependence on aerobic respiration [3]. Roots remain living, energy consuming organs at night, under fluctuating light, and during developmental phases when transport, maintenance, and defence must continue [2]. The available evidence therefore supports a clear conceptual distinction: photosynthesis in the shoot may provide carbon substrates, but oxygen in the root zone is still required for efficient ATP generation within the root itself [1], [3].

V. Practical Implications and Limitations

One practical implication of the evidence is that the common intuition that illuminated plants can somehow dispense with root oxygen is physiologically unsound. A primary study in moss tissues shows that mitochondrial respiration remains essential for ATP supply even in photosynthetically active cells under light [10]. Although this is not a root specific study, it strengthens the broader plant physiological principle that mitochondrial energy metabolism is not simply switched off by photosynthesis. For hydroponics, the implication is cautious but important: if illuminated cells still require mitochondrial respiration for ATP supply, non photosynthetic root tissues are unlikely to be any less dependent on oxygen driven aerobic metabolism [10].

At the same time, the evidence base has limits that should be stated explicitly. The moss study supports the general link between metabolic activity and oxygen dependent ATP generation, but it does not quantify oxygen demand in hydroponic roots or specify thresholds across crops and stages [10]. Likewise, some hydroponic reviews report practical thresholds, noting that dissolved oxygen above 5 mg/L is typically required for effective root uptake and that stagnant water can harm roots [4]. These are useful operational signals, but they do not by themselves explain the mechanism of root respiration in depth. The strongest supported conclusion is

therefore qualitative and physiological, not a universal numeric rule for every hydroponic context.

A further practical lesson is that oxygen management should be treated as a moving target influenced by system conditions and plant demand. Additional reviews note that deep water and floating systems require aeration to avoid oxygen deficiency, and that stagnant conditions can become anaerobic and harmful to roots, while elevated temperatures can both reduce dissolved oxygen availability and increase plant oxygen demand [5], [11], [12]. These claims support the need for continuous root oxygenation, but they remain brief and largely management oriented rather than mechanistically detailed. Future validation would therefore be strongest where root respiration, dissolved oxygen, crop stage, and temperature are examined together within specific hydroponic systems [4], [12].

VI. Conclusion

The evidence supports a straightforward answer to the central question: hydroponic roots need oxygen because they respire continuously to generate ATP through aerobic respiration, and that energy is required for maintenance, nutrient uptake, hydraulic regulation, growth, and defence [1], [2]. Light exposure above the root zone does not remove this requirement, because photosynthesis in shoots does not substitute for oxygen dependent mitochondrial ATP production within root tissues themselves [1], [10]. Hydroponic system design matters chiefly because it can either sustain or constrain the delivery of oxygen to a biologically active root environment [3], [8].

The main limitation of the current evidence is not whether root oxygen matters, but how precisely oxygen demand varies across crops, growth stages, temperatures, and hydroponic configurations. The supplied literature repeatedly recommends oxygenation and, in some cases, reports indicative dissolved oxygen values, yet much of it remains stronger on management guidance than on root specific respiratory quantification [4], [5], [9], [12]. Even with that limitation, the supported conclusion is robust: in hydroponics, root zone oxygen is a continuous biological requirement rather than an optional system feature.

Acknowledgment

The author used artificial intelligence tools to assist with manuscript drafting and revision, and with LaTeX and bibliography preparation. The author reviewed all AI-assisted output and remains responsible for the final content.

References

[1] J. Jethva, R. R. Schmidt, M. Sauter, and J. Selinski, "Try or die: Dynamics of plant respiration and how to survive low oxygen conditions," *Plants*, 2022.

[2] S. Scott, "Dissolved oxygen limitation and pythium root rot in strawberry nft systems: mechanisms, research gaps, and prospects for substrate-free production," *Scott S and Villouta C (*, 2026.

[3] I. A. Lakhiar, H. Yan, T. N. Syed, C. Zhang, S. A. Shaikh, M. Rakibuzzaman, and R. B. Vistro, "Soilless agricultural systems: Opportunities, challenges, and applications for enhancing horticultural resilience to climate change and urbanization," *Horticulturae*, 2025.

[4] M. Elakrouch, M. Mohaddab, A. Rachid, and M. H. Jijakli, "Soilless agriculture at a crossroads: Strengths, challenges, and prospects of hydroponics, aquaponics, and bioponics in relation to precision farming," *Sustainability*, 2026.

[5] J. Arshad, F. Azeem, A. Butt, M. Chaudhary, R. S. Safdar, M. K. Joyo, I. Ahmad, P. Valsalan, and H. M. Ahmed, "Ai-enabled iot-based hydroponic farming with embedded automation and nutrient prediction," *Future Internet*, 2026.

[6] S. Wang, Y. Kleiner, S. M. Clark, V. Raghavan, and B. Tartakovsky, "Review of current hydroponic food production practices and the potential role of bioelectrochemical systems," *Reviews in Environmental Science and Bio/Technology*, 2024.

[7] K. Vought, H. K. Bayabil, J. Pompeo, D. Crawford, Y. Zhang, M. Correll, and A. Martin-Ryals, "Dynamics of micro and macronutrients in a hydroponic nutrient film technique system under lettuce cultivation," *Heliyon*, 2024.

[8] A. Tuxun, Y. Xiang, Y. Shao, J. E. Son, M. Yamada, S. Yamada, K. Tagawa, B. Baiyin, and Q. Yang, "Soilless cultivation: Precise nutrient provision and growth environment regulation under different substrates," *Plants*, 2025.

[9] R. S. Velazquez-Gonzalez, A. L. Garcia-Garcia, J. O. Barceinas-Sanchez, E. Ventura-Zapata, and J. C. Sosa-Savedra, "A review on hydroponics and the technologies associated for medium- and small-scale operations," *Agriculture*, 2022.

[10] A. M. Vera-Vives, P. Novel, K. Zheng, S. ling Tan, and M. Schwarzländer, "Mitochondrial respiration is essential for photosynthesis-dependent atp supply of the plant cytosol," *New Phytologist*, 2024.

[11] C. Mai, A. Mojiri, S. Palanisami, A. Altaee, Y. Huang, and J. L. Zhou, "Wastewater hydroponics for pollutant removal and food production: Principles, progress and future outlook," *Water*, 2023.

[12] A. Fussy and J. Papenbrock, "An overview of soil and soilless cultivation techniques-chances, challenges and the neglected question of sustainability," *Plants*, 2022.