Dissolved oxygen and water temperature

Grant Kinkead
Independent Researcher, Melbourne, VIC, Australia
grant@kinkead.au

***Abstract*—Dissolved oxygen is a central constraint in hydroponic root-zone management because oxygen supply to roots depends on both the physical capacity of water to hold oxygen and the biological demand imposed by plants and associated organisms. This paper explains why water temperature changes the oxygen available to hydroponic roots by synthesising evidence from recent hydroponic and water oxygenation literature. The reviewed evidence shows a consistent physical trend: as water temperature rises, equilibrium oxygen solubility falls, with modelling and experimental data showing lower saturation concentrations at 30 degrees Celsius than at 24 degrees Celsius [1]. However, dissolved oxygen concentration alone does not define root risk. Root oxygen demand is a separate biological variable that can increase during active growth and under stressful root-zone conditions, so warmer solution can reduce oxygen supply while also intensifying demand and disease pressure [2], [3]. Across hydroponic reviews and system studies, operational dissolved oxygen targets are often discussed, yet the distinction between concentration and demand is not always made explicit [4], [5]. The main implication is that temperature management is not merely a comfort setting for roots, but a governing factor in root-zone oxygen balance, monitoring strategy, and risk control in hydroponic production [6], [7].**

I. Introduction

In hydroponic systems, dissolved oxygen and water temperature are tightly linked because the amount of oxygen that water can hold at equilibrium is temperature dependent. Recent thermodynamic and experimental work on oxygenation shows that oxygen solubility decreases as water temperature increases, with reported saturation concentrations lower at 30 degrees Celsius than at 24 degrees Celsius, a pattern consistent with Henry's law based interpretation of gas dissolution in water [1]. For hydroponic root zones, this means that warming the nutrient solution changes the physical ceiling on available oxygen before any root or microbial consumption is considered. The immediate implication is that growers cannot interpret a warm reservoir and a cool reservoir using the same oxygen margin, even when aeration hardware appears unchanged [1].

A second issue is that oxygen availability to roots cannot be understood from dissolved oxygen concentration alone. Evidence from a review of strawberry nutrient film technique systems distinguishes between the amount of oxygen present in solution and the respiratory demand imposed by roots at different developmental stages. In that literature, passive oxygen supply can become inadequate while root respiratory demand rises during establishment and fruiting, showing that concentration and demand are separate variables that interact in practice [2]. This distinction matters because a reading that appears acceptable on a sensor may still coincide with root stress if demand has increased, making interpretation of dissolved oxygen inherently biological as well as physical [2].

Temperature therefore influences hydroponic oxygen risk through two coupled pathways rather than one. On the supply side, warmer water holds less oxygen at equilibrium. On the demand side, the biological system can become more oxygen intensive, while stressful warm conditions may also favour root dysfunction and disease processes that further destabilise the rhizosphere [2]. This paper develops that argument as an evidence based explanation rather than a new experiment. It first defines the topic and its scope, then explains the governing mechanisms, synthesises the available hydroponic evidence, and finally identifies the practical implications and remaining limits of current knowledge for hydroponic root-zone management [1], [2].

II. Background and Scope

Existing hydroponic literature often recognises that cooler nutrient solution can improve oxygen conditions, but the emphasis is not always on the underlying physical explanation. The strongest review evidence in the present source set reports that reducing nutrient-solution temperature towards about 18 to 22 degrees Celsius can increase oxygen solubility by roughly 15 to 20 percent in hydroponic context [2]. That observation clearly establishes the direction of the temperature effect required for this paper: cooler water generally supports higher equilibrium dissolved oxygen than warmer water. Even so, the review is primarily concerned with oxygen limitation and root disease in strawberry nutrient film systems, so its main contribution here is to confirm the practical importance of the temperature effect rather than to provide a full general theory of gas solubility in hydroponics [2].

Broader reviews of soilless cultivation also reinforce the importance of root-zone temperature control, although they usually treat it as part of general environmental management rather than as a dedicated oxygen balance problem. One review recommends nutrient-solution temperatures around 18 to 22 degrees Celsius and emphasises controlled root-zone conditions as part of resilient soilless production [6]. This is valuable contextual support because it shows that temperature management is already considered operationally important. However, that literature does not by itself explain the two-part mechanism requested here. It gives useful agronomic guidance, but it does not fully separate the physical reduction in oxygen solubility from the distinct biological question of how much oxygen roots and related organisms require under changing conditions [6].

The most direct contribution on the physical side comes from the thermodynamic oxygenation study, which quantifies dissolved oxygen behaviour under different temperatures and oxygenation conditions [1]. That work helps clarify why temperature matters to concentration by showing both modelled and experimental differences in oxygen saturation and transfer performance. Yet it is not a hydroponic root physiology paper, and it does not explain root oxygen demand as a separate biological variable [1]. The gap that remains, and that defines the scope of this paper, is therefore interpretive rather than empirical: available evidence supports the temperature effect on dissolved oxygen concentration and separately supports the importance of root demand, but the literature must be synthesised to explain how these processes combine into root-zone oxygen risk in hydroponics [1], [2].

III. Core Concepts and Mechanisms

The first core mechanism is physical solubility. Oxygen enters and remains in water according to thermodynamic constraints, so warmer water has a lower equilibrium capacity for dissolved oxygen than cooler water. The oxygenation study in the present evidence base supports this directly, reporting lower saturation concentration at higher temperature and showing that oxygenation efficiency also declines as temperature rises [1]. For hydroponics, this means that water temperature changes not only the final saturation point but also the operating ease with which that point can be approached. In practical terms, a system may need more active aeration or more favourable flow conditions to maintain a similar dissolved oxygen reading in warmer solution, and even then the physical ceiling remains lower [1].

The second mechanism is the distinction between concentration and demand. Hydroponic management discussions often specify dissolved oxygen targets because solution concentration is measurable and operationally convenient. Reviews of soilless agriculture note the importance of dissolved oxygen in hydroponic and aquaponic operation, indicating that concentration is treated as a meaningful control variable [4]. However, concentration is only the supply state of the water, not the whole oxygen economy of the root zone. Roots must consume oxygen for respiration, and the significance of any measured concentration depends on whether that supply is adequate for the current biological load. The implication is that a target value is informative but incomplete unless it is interpreted alongside crop stage, temperature regime, and the broader root-zone context [4].

The third mechanism concerns the interface between measured solution oxygen and actual root need. A prototype hydroponic system study reports maintaining dissolved oxygen above 5 mg/L for root respiration in a deep-water culture subsystem, linking the metric directly to biological function [5]. That connection is useful because it shows why dissolved oxygen readings are not merely chemical descriptors; they are operational proxies for respiratory support. At the same time, the study does not clearly disentangle whether an acceptable concentration under one temperature and crop condition would remain acceptable under another [5]. This limitation underscores the central argument of the present paper: dissolved oxygen concentration is necessary to monitor, but its meaning depends on how temperature reshapes both the supply side of oxygen dissolution and the demand context within the root zone [1], [5].

IV. Evidence Synthesis

When the accepted sources are read together, they support a layered interpretation of oxygen risk rather than a single threshold rule. The hydroponic system paper provides an operational framing by linking dissolved oxygen targets with root respiration needs, but it does not explain how temperature shifts the risk structure behind those targets [5]. The strawberry nutrient film review fills that interpretive gap by showing that oxygen limitation can emerge where passive oxygen supply becomes inadequate at the same time that root respiratory demand increases during critical growth phases [2]. Taken together, these sources suggest that oxygen management in hydroponics should not be treated as maintaining one fixed number in solution. Instead, risk depends on whether the measured concentration remains sufficient after temperature has already reduced the physical oxygen reserve and biological activity has increased consumption pressure [2], [5].

A similar pattern appears in nutrient-solution management literature. Reviews of hydroponic nutrient management identify dissolved oxygen monitoring and climate related variables as important parts of system performance, indicating that oxygen status is embedded within broader environmental control rather than isolated from it [7]. This supports the idea that temperature must be interpreted as a system variable with consequences for oxygen availability, not simply as a separate environmental setting. Yet these reviews do not directly articulate the two-pathway model of risk in which warmer water both reduces oxygen solubility and alters biological demand [7]. The main synthesis point is therefore analytical: hydroponic practice already recognises oxygen and temperature as linked management concerns, but the evidence becomes more explanatory when physical solubility studies and root-zone biological studies are considered together [1], [7].

The broadest conceptual confirmation comes from review literature noting that high temperatures can reduce dissolved oxygen while increasing plant oxygen demand [3]. This statement is important because it captures the reader-facing answer in compact form, but it is less developed mechanistically than the strawberry review and less quantitative on the physical side than the thermodynamic oxygenation study [1], [2], [3]. Even so, the consistency across these sources strengthens the overall conclusion. There is no serious conflict in direction: warmer water reduces the equilibrium oxygen supply, and warmer root-zone conditions can intensify the biological context in which oxygen is consumed. The remaining limitations concern depth of explanation, crop specificity, and the degree to which general hydroponic recommendations can be translated into crop specific oxygen risk assessment [2], [3].

V. Practical Implications and Limitations

For hydroponic practice, the main implication is that water temperature management is inseparable from oxygen management. If dissolved oxygen targets are monitored without considering temperature, operators may misjudge how much oxygen security is actually present in the root zone [5], [7]. A concentration that appears satisfactory under cooler conditions may represent a much narrower safety margin under warmer conditions because the equilibrium supply ceiling has fallen and the biological system may be consuming oxygen more rapidly [3]. This does not justify a universal numeric rule from the present evidence, because the available sources differ in system type and analytical purpose. It does, however, justify a robust operational principle: dissolved oxygen readings should be interpreted in relation to temperature, crop condition, and root-zone stress context rather than as free-standing indicators [3], [5], [7].

The evidence base also has clear limitations that should shape interpretation. Several accepted sources are reviews or prototype oriented studies rather than comparative hydroponic experiments designed specifically to isolate temperature, dissolved oxygen concentration, and root oxygen demand as separate variables [5], [7]. The thermodynamic study offers strong support for the physical direction of the solubility effect, but it is not a direct study of hydroponic roots [1]. Conversely, hydroponic reviews often identify appropriate temperature ranges or oxygen concerns, yet do not provide a full mechanistic partition between concentration and demand [3], [6]. The highest-value validation need is therefore integrated hydroponic research that simultaneously measures water temperature, dissolved oxygen concentration, crop stage, and indicators of root oxygen demand or stress, so that oxygen risk can be interpreted as a balance rather than as a single reservoir measurement [2], [7].

VI. Conclusion

The evidence supports a clear answer to the question of why water temperature changes the oxygen available to hydroponic roots. Warmer water generally holds less dissolved oxygen at equilibrium, so the physical supply potential of the nutrient solution declines as temperature rises [1]. At the same time, dissolved oxygen concentration is not identical to root oxygen demand. Hydroponic root zones can become more vulnerable when biological demand increases during active growth or stressful conditions, meaning that temperature changes the demand context as well as the supply ceiling [2], [3]. Across the accepted literature, this dual interpretation is more scientifically defensible than any single-threshold view of oxygen sufficiency [4], [5]. The most important next step is better integrated hydroponic evidence that links temperature, dissolved oxygen readings, and root respiratory stress in the same experimental frame, but the present synthesis already establishes that temperature is a governing variable in root-zone oxygen balance, not a secondary environmental detail [6], [7].

Acknowledgment

The author used artificial intelligence tools to assist with manuscript drafting and revision, and with LaTeX and bibliography preparation. The author reviewed all AI-assisted output and remains responsible for the final content.

References

[1] C. D. Mihaela Constantin and M. Oprea, "Thermodynamic analysis of oxygenation methods for stationary water: Mathematical modeling and experimental investigation," *Thermo*, 2025.

[2] S. Scott, "Dissolved oxygen limitation and pythium root rot in strawberry nft systems: mechanisms, research gaps, and prospects for substrate-free production," *Scott S and Villouta C (*, 2026.

[3] A. Fussy and J. Papenbrock, "An overview of soil and soilless cultivation techniques-chances, challenges and the neglected question of sustainability," *Plants*, 2022.

[4] M. Elakrouch, M. Mohaddab, A. Rachid, and M. H. Jijakli, "Soilless agriculture at a crossroads: Strengths, challenges, and prospects of hydroponics, aquaponics, and bioponics in relation to precision farming," *Sustainability*, 2026.

[5] J. Arshad, F. Azeem, A. Butt, M. Chaudhary, R. S. Safdar, M. K. Joyo, I. Ahmad, P. Valsalan, and H. M. Ahmed, "Ai-enabled iot-based hydroponic farming with embedded automation and nutrient prediction," *Future Internet*, 2026.

[6] I. A. Lakhiar, H. Yan, T. N. Syed, C. Zhang, S. A. Shaikh, M. Rakibuzzaman, and R. B. Vistro, "Soilless agricultural systems: Opportunities, challenges, and applications for enhancing horticultural resilience to climate change and urbanization," *Horticulturae*, 2025.

[7] E. Fathidarehnijeh, M. Nadeem, M. Cheema, R. Thomas, M. Krishnapillai, and L. Galagedara, "Current perspective on nutrient solution management strategies to improve the nutrient and water use efficiency in hydroponic systems," *Canadian Journal of Plant Science*, 2024.